

# **NuFact 2018, 20th workshop on neutrinos from accelerators**

Sunday 12 August 2018 - Saturday 18 August 2018

## **Book of Abstracts**



# Contents

A High-efficiency Cosmic Ray Veto Detector for the Mu2e Experiment . . . . .	1
A reappraisal of constraints on Z-prime models from unitarity and direct searches at the LHC . . . . .	1
A short travel for neutrinos in Large Extra Dimensions . . . . .	1
ANNIE Phase II Detector and Event Reconstruction . . . . .	1
Ab Initio Methods . . . . .	2
Atmospheric Neutrino Oscillations with IceCube/DeepCore . . . . .	2
Atmospheric Neutrino Oscillations with IceCube/DeepCore . . . . .	2
Atmospheric neutrino results from Super-Kamiokande . . . . .	2
Atmospheric neutrino results from Super-Kamiokande . . . . .	2
Average CsI neutron density distribution from COHERENT data . . . . .	2
BELLEII (TBC) . . . . .	2
BSM Neutrino Theory . . . . .	3
Boosted Dark Matter at DUNE . . . . .	3
CAPTAIN Results . . . . .	3
Close . . . . .	3
Closing Talk . . . . .	3
Coherent Elastic Neutrino Nucleus Scattering (CEvNS) as a probe of Z-prime through kinetic and mass mixing effects . . . . .	3
Coherent Scattering Results and Future . . . . .	3
Cold muonium beam for atomics physics and gravity experiments . . . . .	4
Commissioning and first results of the Fermilab Muon Campus . . . . .	4
Constraining neutrino transition magnetic moments . . . . .	4
Constraints on neutrino decay scenarios with electron anti-neutrino disappearance experiments. . . . .	4

Cross section issues for the next decade . . . . .	5
DC Muon Beam Physics . . . . .	5
DUNE Near Detector Plans . . . . .	5
DUNE Oscillation Physics . . . . .	5
DUNE and CPT-violating Neutrinos . . . . .	5
Data Analysis and Preliminary Results of the Proton Charge Radius Experiment at JLab . . . . .	6
Daya Bay Reactor Neutrino Experiment . . . . .	6
Design and Challenges of ESSnusb Accumulator . . . . .	6
Details of the NOvA oscillation analyses . . . . .	6
Details of the T2K oscillation analyses . . . . .	6
Details of the T2K oscillation analyses . . . . .	7
Detector concepts for nuSTORM . . . . .	7
Development and operational experience of T2K magnetic horn for over-MW beam . . . . .	7
Discussion . . . . .	7
Discussion . . . . .	7
Discussion . . . . .	7
Discussion . . . . .	8
Discussion: (re)forming a nuSTORM collaboration . . . . .	8
Distinguishing muon LFV effective couplings using $\mu \rightarrow e e \gamma$ . . . . .	8
Distinguishing muon LFV effective couplings using $\mu \rightarrow e e \gamma$ . . . . .	8
Double Chooz . . . . .	8
EMuS at CSNS Updated Studies . . . . .	8
ESSnusb Target and Horn Studies and Future Development . . . . .	9
Electron Scattering . . . . .	9
Electron scattering data and neutrinos . . . . .	9
Electron vs Muon Neutrinos . . . . .	9
Exploring the Potential of Short-Baseline Physics at Fermilab . . . . .	9
FNAL SBL Program Status . . . . .	9
FNAL SBN Status . . . . .	9
Fabrication of A High-efficiency Cosmic Ray Veto Detector for the Mu2e Experiment . . . . .	10

Final results from the OPERA experiment in the CNGS neutrino beam . . . . .	10
Final results from the OPERA experiment in the CNGS neutrino beam . . . . .	10
Final results from the OPERA experiment in the CNGS neutrino beam . . . . .	10
First Results from the PROSPECT Short Baseline Reactor Experiment . . . . .	11
Future DUNE constraints on EFT . . . . .	11
Future Upgrade of J-PARC Target and Beam Window . . . . .	11
Future prospects for CAPTAIN experiment . . . . .	11
GENIE Physics Tuning . . . . .	12
Global Neutrino Oscillation Fits . . . . .	12
Global analysis of neutrino oscillation experiments . . . . .	12
Integrable Optics Test Accelerator . . . . .	12
IsoDAR . . . . .	12
LArIAT Cross Section Results . . . . .	12
LHC Results on heavy Neutrino Searches . . . . .	12
Latest Results from MINOS+ on Sterile Neutrinos search . . . . .	13
Latest Results from the Daya Bay Reactor Neutrino Experiment . . . . .	13
Lattice QCD and neutrinos . . . . .	13
Lepton Flavour Universal at LHCb . . . . .	13
Lepton-Number-Charged Scalars and Neutrino Beamstrahlung . . . . .	13
Light scalar dark matter at neutrino oscillation experiments . . . . .	13
Low Emittance Muon Accelerator . . . . .	13
Low Emittance Muon Beams . . . . .	14
Low energy neutrino interactions . . . . .	14
MICE Results . . . . .	14
MINERvA Cross Section Results . . . . .	14
Measurement of Phase-Space Density Evolution in MICE . . . . .	14
Measuring the Leptonic Dirac CP Phase with Muon Decay at Rest . . . . .	14
MicroBooNE Cross Section Results . . . . .	14
MicroBooNE Search for Low-Energy Excess Using Deep Learning Algorithms . . . . .	15
Mini-CAPTAIN measurements in the LANSCE WNR Neutron Beam . . . . .	15

Model independent non-unitarity . . . . .	15
Modeling neutrino-nucleus interactions in the few-GeV region . . . . .	15
Mu2e II - A Proposed Evolution of the Mu2e Experiment . . . . .	16
Muonic X-ray measurements with radioactive elements . . . . .	16
NOvA Cross Section Model / Oscillation Needs . . . . .	16
NOvA Cross Section Results . . . . .	16
NUISANCE for neutrino cross section fits . . . . .	17
Neutrino Dark Matter Connections . . . . .	17
Neutrino Oscillations in Dark Backgrounds . . . . .	17
Neutrino Physics with Deep Learning on NOvA . . . . .	17
Neutrino Physics with Deep Learning on NOvA . . . . .	17
Neutrino flavor transformation in supernova as a probe for nonstandard neutrino-scalar interactions . . . . .	18
Neutrino physics with KM3NeT/ORCA . . . . .	18
Neutrino physics with KM3NeT/ORCA . . . . .	18
Neutrino trident at DUNE . . . . .	18
Neutrino trident production at near detectors . . . . .	18
New Results from RENO . . . . .	19
Nuclear Theory, Data and Event Generators . . . . .	19
Nucleon axial form-factor from a Bayesian neural network analysis of scattering data . . . . .	19
Open Issues in Nuclear SIS and DIS Scattering . . . . .	19
Overview of Neutrino Physics . . . . .	19
Performance of Scintillation Counters as Measured at the Fermilab Test Beam Facility for the Mu2e Cosmic Ray Veto System . . . . .	20
Performance of Wavelength-Shifting Fibers for the Mu2e Cosmic Ray Veto Detector . . . . .	20
Physics potential of Hyper-Kamiokande for neutrino oscillation measurements . . . . .	20
Physics potential of the ESSvSB facility . . . . .	20
Precision spectroscopy of exotic atoms involving muon . . . . .	21
Probing Neutrino Mass Ordering and Solar neutrinos with JUNO detector . . . . .	21
Probing neutrino coupling to a light scalar with coherent neutrino scattering . . . . .	21

Probing secret interactions of eV-scale sterile neutrinos with the diffuse supernova neutrino background . . . . .	21
Pulsed Muon Beam Physics . . . . .	21
Reactor Antineutrino Detection Using CHANDLER : A New Portable Neutrino Detector Tulasi Subedi Abstract CHANDLER . . . . .	21
Recent Cross Section Results from T2K . . . . .	22
Recent Progress on Radiation Damage Studies at RaDIATE . . . . .	22
Recent Results from MICE on Multiple Coulomb Scattering and Energy Loss . . . . .	22
Recent Results from MINERVA . . . . .	22
Recent Results from the Study of Emittance Evolution in MICE . . . . .	23
Recent Results from the T2K Near Detector . . . . .	23
Recent progress on radiation damage studies at RaDIATE . . . . .	23
Recent results of neutrino interactions from the T2K Near Detector . . . . .	23
Removal and Binding Energy in Lepton-Nucleus Scattering . . . . .	24
Results and Prospects from NOvA . . . . .	24
Results and Prospects from T2K . . . . .	24
Results from IceCube . . . . .	24
Review of Tension in Data/Models of Neutrino Cross Sections . . . . .	24
SBL Reactor Experiments . . . . .	24
Search for $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ at CERN . . . . .	25
Search for lepton favour violation with the ATLAS detector . . . . .	25
Searches for Electric Dipole Moments (EDM) at a Storage Ring with JEDI . . . . .	25
Searches for heavy neutral lepton production and lepton flavour violation in kaon decays at the NA62 experiment . . . . .	25
Shedding light on low energy excess anomaly with MiniBooNE and MicroBooNE . . . . .	25
Simulation studies of a detector for nuSTORM . . . . .	25
Single Ion Barium Tagging for Neutrino-less Double-Beta Decay Searches. A multi-disciplinary technique in development for the NEXT experiment . . . . .	26
Sketch of physics case in preparation for the EU Strategy Update . . . . .	26
Solar $\nu$ at DUNE and BSM . . . . .	26
Standard Model prediction for the muon $g-2$ . . . . .	26

Standard and non-standard neutrino physics at reactor experiments . . . . .	26
Status and Future of high-power Neutrino Target Stations . . . . .	26
Status and Future of high-power Proton Drivers . . . . .	27
Status and Physics Potential of MOMENT Study . . . . .	27
Status and Physics of JUNO . . . . .	27
Status of Charged Lepton Flavor Violation searches at CMS and future prospects . . . . .	27
Status of DUNE . . . . .	27
Status of ESSnuB . . . . .	27
Status of NA61/SHINE Measurements for Neutrino Experiments . . . . .	27
Status of ProtoDUNE Experiments at CERN . . . . .	28
Status of T2HK . . . . .	28
Status of accelerator-based Neutrino Physics . . . . .	28
Status of consideration of implementation of nuSTORM at CERN . . . . .	28
Status of the COMET Experiment . . . . .	28
Status of the DeeMe Experiment . . . . .	28
Status of the ENUBET . . . . .	29
Status of the Facility/Accelerator/Beam-line for Muon Programs at J-PARC . . . . .	29
Status of the ISODAR project . . . . .	29
Status of the MEG II Experiment . . . . .	29
Sterile Neutrinos search via NC dis at NOvA . . . . .	29
Sterile neutrino searches with the ICARUS detector . . . . .	29
Sterile neutrinos searches . . . . .	29
Studies of PRISM/PRIME - the next generation muon to electron conversion experiment	30
Studies of the Aging Properties of the Mu2e Cosmic Ray Veto System . . . . .	30
Studying Neutral Current Elastic Scattering and the Strange Axial Form Factor in Micro-BooNE . . . . .	30
T2K Cross Section Model / Oscillation Needs . . . . .	31
T2K Cross Section Results . . . . .	31
T2K Near Detector Upgrades and Plans for T2HK . . . . .	31
Test on lepton flavor universality at BESIII . . . . .	31



The LBNF Beamline . . . . .	31
The MUon Scattering Experiment (MUSE) at the Paul Scherrer Institute . . . . .	31
The Mu2e Experiment at Fermilab . . . . .	31
The Mu2e Experiment at Fermilab . . . . .	32
The NOvA Test Beam Program . . . . .	32
The Role of Cross Sections in the Oscillation Analysis: The T2K Experience . . . . .	32
The design and research progresses of the Central Detector in JUNO . . . . .	33
The g-2 Beamline . . . . .	33
The g-2 experiment at Fermilab . . . . .	33
The search for Lepton-Flavour Violation with the Mu3e Experiment . . . . .	33
The search for neutral currents in muonic X-rays . . . . .	33
Theory of electron scattering and neutrinos . . . . .	34
Theory of neutrino pion production . . . . .	34
Towards a new High Intensity Muon Beam at PSI: Status and Prospects . . . . .	34
Upgrade Possibility of the ESS Linac for the ESSnuSB Project . . . . .	34
WAGASCI . . . . .	34
WG1 Goals for the Meeting . . . . .	35
WG1 Summary . . . . .	35
WG2 Goals for the Meeting . . . . .	35
WG2 Summary . . . . .	35
WG3 Goals for the Meeting . . . . .	35
WG3 Summary . . . . .	35
WG4 Charge . . . . .	35
WG4 Goals for the Meeting . . . . .	36
WG4 Summary . . . . .	36
WG4 and WG5 - Working Time . . . . .	36
WG5 Goals for the Meeting . . . . .	36
WG5 Summary . . . . .	36
Welcome . . . . .	36
Welcome . . . . .	36

Welcome and introduction . . . . .	37
Working Group Discussion Time . . . . .	37
Working Group Discussion Time . . . . .	37
nuSTORM Facility and Report from Cross Section Workshop . . . . .	37
nuSTORM accelerator concept to serve cross-section programme . . . . .	37

**Posters & welcome reception / 48****A High-efficiency Cosmic Ray Veto Detector for the Mu2e Experiment****Author(s):** Ben Barton<sup>1</sup> ; Danny Mills<sup>1</sup>**Co-author(s):** Craig Group<sup>2</sup><sup>1</sup> *Virginia*<sup>2</sup> *University of Virginia***Corresponding Author(s):** dm8bz@virginia.edu, group@virginia.edu, bb6yx@virginia.edu

The Mu2e experiment at Fermilab will search for the charged lepton flavor violating process of coherent muon-to-electron conversion in the presence of a nucleus with a sensitivity four orders of magnitude beyond current limits. The experiment will have a single event sensitivity of about  $3 \times 10^{-17}$  while limiting the total background to about 0.5 events. One potential background is due to cosmic-ray muons producing an electron that is indistinguishable from signal within the Mu2e apparatus. The cosmic-ray-veto system of the Mu2e experiment is tasked with vetoing cosmic-ray-induced backgrounds with high efficiency, without inducing significant dead time and while operating in a high-intensity environment. The design of the cosmic-ray-veto system will be discussed.

**WG5 / 91****A reappraisal of constraints on Z-prime models from unitarity and direct searches at the LHC****Corresponding Author(s):** gondogolegogol@gmail.com**WG5 / 83****A short travel for neutrinos in Large Extra Dimensions****Corresponding Author(s):** gabivstenico@gmail.com**Posters & welcome reception / 204****ANNIE Phase II Detector and Event Reconstruction**Jingbo Wang<sup>None</sup>**Corresponding Author(s):** jiowang@ucdavis.edu

The Accelerator Neutrino Neutron Interaction Experiment (ANNIE), deployed on the Booster Neutrino Beam (BNB) at Fermilab, has recently finished the neutron background measurement in the Phase I data taking. The primary physics goal of Phase II is to measure the multiplicity of final state neutrons from neutrino-nucleus interactions in water, which provides a strong handle to study the systematic uncertainties relevant to the neutrino energy reconstruction in the future long baseline oscillation experiments. The ANNIE Phase II detector will use Gadolinium-loaded water to detect the final state neutrons from neutrino interactions. It will also incorporate five Large Area Picosecond PhotoDetectors (LAPPDs) to improve the vertex and track reconstruction capability required

by the physics goals. This presentation will give an overview of the Phase II detector upgrade and focus on the event reconstruction capability improved by the LAPPDs.

**WG2 / 68**

## **Ab Initio Methods**

**Corresponding Author(s):** ale.lovato85@gmail.com

**WG1 Neutrino oscillation / 179**

## **Atmospheric Neutrino Oscillations with IceCube/DeepCore**

**Corresponding Author(s):** cowen@phys.psu.edu

**WG1 / 170**

## **Atmospheric Neutrino Oscillations with IceCube/DeepCore**

**Corresponding Author(s):** cowen@phys.psu.edu

**WG1 Neutrino oscillation / 178**

## **Atmospheric neutrino results from Super-Kamiokande**

**Corresponding Author(s):** cbronner@km.icrr.u-tokyo.ac.jp

**WG1 / 169**

## **Atmospheric neutrino results from Super-Kamiokande**

**Corresponding Author(s):** cbronner@km.icrr.u-tokyo.ac.jp

**WG5 / 82**

## **Average CsI neutron density distribution from COHERENT data**

**Corresponding Author(s):** matteo.cadeddu@ca.infn.it

**WG4 - CLFV at Colliders / 116**

## **BELLEII (TBC)**

**Plenary X / 36**

## **BSM Neutrino Theory**

**WG5 / 87**

## **Boosted Dark Matter at DUNE**

**Corresponding Author(s):** doojin.kim@cern.ch

**WG1+WG2 / 130**

## **CAPTAIN Results**

**Corresponding Author(s):** jchaves@sas.upenn.edu

**Plenary Summary II and Closeout / 208**

## **Close**

**Corresponding Author(s):** pahuber@vt.edu

**Plenary Summary II and Closeout / 39**

## **Closing Talk**

**Corresponding Author(s):** alexfr@slac.stanford.edu

**WG5 / 84**

## **Coherent Elastic Neutrino Nucleus Scattering (CEvNS) as a probe of Z-prime through kinetic and mass mixing effects**

**Corresponding Author(s):** jbdent@louisiana.edu

Plenary X / 16

## Coherent Scattering Results and Future

Corresponding Author(s): psbarbeau@phy.duke.edu

WG3 and WG4 - Muon Beam Facilities / 127

## Cold muonium beam for atomics physics and gravity experiments

Corresponding Author(s): anna.soter@psi.ch

WG3 and WG4 - Muon Beam Facilities / 128

## Commissioning and first results of the Fermilab Muon Campus

Corresponding Author(s): stratakis@gmail.com

Posters & welcome reception / 58

## Constraining neutrino transition magnetic moments

Nitali Dash<sup>1</sup> ; Reetanjali Moharana<sup>2</sup> ; Guofu Cao<sup>1</sup>

<sup>1</sup> *Institute of Hienergy Physics, Chinese Academy of Sciences, Beijing 100049, China*

<sup>2</sup> *The Racha Institute of Physics, 91904 Jerusalem, Israel*

Corresponding Author(s): dnitali@ihep.ac.cn

We are presenting a preliminary results on the studies of neutrino transition magnetic moments using DUNE LAr, HK and JUNO detectors. Neutrinos, if Majorana particles, the combined effect of magnetic field and matter effect in core-collapse Super Nova can transform some of  $\nu_e$  to  $\bar{\nu}_e$  due to spin flavour conversions. As a result of this conversions the inverse beta decay signal will have an increment indicating evidence of transition magnetic moments. The DUNE LAr is sensitive to  $\nu_e$  so will observed a deficiency of  $\nu_e$  due to this conversion whereas both HK and JUNO which are sensitive to  $\bar{\nu}_e$  will see excess of  $\bar{\nu}_e$ . The DUNE LAr and JUNO are more or less sensitive to other type of neutrinos due to use of  $^{40}\text{Ar}$  and  $^{12}\text{C}$ . So can estimate the event ratio using both neutrinos and hence sensitivity on transition magnetic moments. Even an non observation of such conversion put a restrictive bounds on the neutrino transition magnetic moments.

Posters & welcome reception / 60

## Constraints on neutrino decay scenarios with electron anti-neutrino disappearance experiments.

Author(s): Suprabh Prakash<sup>1</sup>

**Co-author(s):** Yago Phillipe Porto Silva <sup>2</sup> ; Orlando L. G. Peres <sup>3</sup> ; Hiroshi Nunokawa <sup>4</sup> ; Hisakazu Minakata <sup>5</sup>

<sup>1</sup> IFGW, Unicamp

<sup>2</sup> Instituto de Física Gleb Watagin

<sup>3</sup> Instituto de Física Gleb Wataghin

<sup>4</sup> PUC, Rio de Janeiro

<sup>5</sup> IFT, UAM

**Corresponding Author(s):** suprabhprakash@gmail.com, nunokawa@puc-rio.br, yporto@ifi.unicamp.br, orlando@ifi.unicamp.br

Neutrino decay provides a very interesting case for the “beyond PMNS” neutrino physics. It has been shown that this phenomenon can also explain some of the anomalies seen in neutrino experiments. We study the constraints that  $\bar{\nu}_e$  disappearance experiments like JUNO and KamLAND can put on neutrino decay scenarios. In particular, we consider a model where a heavier neutrino can decay giving active daughter neutrinos which can then be detected in these experiments. We find that the experiments JUNO and KamLAND can together constrain  $\tau_3/m_3$   $10^{-10}$  s/eV for the normal hierarchy and  $\tau_2/m_2$   $10^{-9}$  s/eV for the inverted hierarchy. We discuss an interesting physics case because of which the bounds are better for the inverted hierarchy. Unlike  $\nu_e$  appearance experiments, the  $\bar{\nu}_e$  disappearance events do not change much depending on whether the decay products are visible or not. This is due to the smallness of  $|Ue3|$ .

**Session 1 / 197**

## Cross section issues for the next decade

**Corresponding Author(s):** morfin@fnal.gov

**Plenary IX / 32**

## DC Muon Beam Physics

**Corresponding Author(s):** angela.papa@psi.ch

**Plenary VI / 29**

## DUNE Near Detector Plans

**Corresponding Author(s):** deroeck@mail.cern.ch

**WG1 Neutrino oscillation / 186**

## DUNE Oscillation Physics

**Corresponding Author(s):** animesh.chatterjee@uta.edu

**WG5 / 98****DUNE and CPT-violating Neutrinos****Corresponding Author(s):** chternes@ific.uv.es**WG4 - Muonic atoms/proton radius / 123****Data Analysis and Preliminary Results of the Proton Charge Radius Experiment at JLab****Corresponding Author(s):** cg225@phy.duke.edu**Posters & welcome reception / 203****Daya Bay Reactor Neutrino Experiment**Shengchao Li<sup>None</sup>**Corresponding Author(s):** scli@vt.edu

Starting in 2011, the Daya Bay Reactor Neutrino Experiment observed anti-neutrinos from six nuclear reactors with eight identically designed underground anti-neutrino detectors in three experimental halls, and has accumulated the world's largest dataset of anti-neutrino candidates. The measurement of the neutrino mixing angle  $\theta_{13}$  and the neutrino mass squared difference  $|\Delta m^2_{ee}|$  have reached a precision of better than 4%. The large dataset allows study of a variety of topics in neutrino physics, such as absolute reactor flux and spectrum. In this poster, we will present the latest results from Daya Bay on several topics.

**WG3 Parallel Session 4 / 153****Design and Challenges of ESSnusB Accumulator****Corresponding Author(s):** ye.zou@physics.uu.se**WG1 Neutrino oscillation / 182****Details of the NOvA oscillation analyses****Corresponding Author(s):** esmith227@gmail.com**WG1 Neutrino oscillation / 183****Details of the T2K oscillation analyses****Corresponding Author(s):** davide.sgalaberna@cern.ch



**Posters & welcome reception / 57**

## **Details of the T2K oscillation analyses**

Davide Sgalaberna<sup>1</sup>

<sup>1</sup> *CERN*

**Corresponding Author(s):** [davide.sgalaberna@cern.ch](mailto:davide.sgalaberna@cern.ch)

T2K is a long-baseline neutrino experiment in which a muon neutrino beam produced by J-PARC in Tokai is sent 295 km across Japan to the Super-Kamiokande detector. The experiment studies neutrino oscillations via the disappearance of muon neutrinos and the appearance of electron neutrinos. T2K has conclusively observed muon neutrino to electron neutrino oscillations, opening the door to the observation of CP violation in the lepton sector. Since 2014, the experiment has run alternating neutrino and antineutrino beams in order to precisely measure the corresponding oscillation probabilities, resulting in leading measurements of the muon antineutrino disappearance parameters and results on CP violation in the lepton sector. Different oscillation analyses are performed. They differ for the adopted statistical approach, either frequentist or bayesian, and the kinematical variables used for the analysis templates. In this talk, we will present recently-updated results, focusing on the details of the oscillation analysis methods.

**Session 1 / 198**

## **Detector concepts for nuSTORM**

**Corresponding Author(s):** [paul.soler@glasgow.ac.uk](mailto:paul.soler@glasgow.ac.uk)

**WG3 Parallel Session 4 / 152**

## **Development and operational experience of T2K magnetic horn for over-MW beam**

**Corresponding Author(s):** [tetsuro.sekiguchi@kek.jp](mailto:tetsuro.sekiguchi@kek.jp)

**WG3 Parallel Session 7 / 157**

## **Discussion**

**Corresponding Author(s):** [tetsuro.sekiguchi@kek.jp](mailto:tetsuro.sekiguchi@kek.jp), [bfreemire@niu.edu](mailto:bfreemire@niu.edu)

**WG3 Parallel Session 3 / 151**

## **Discussion**

**Corresponding Author(s):** [bfreemire@niu.edu](mailto:bfreemire@niu.edu), [tetsuro.sekiguchi@kek.jp](mailto:tetsuro.sekiguchi@kek.jp)

**WG2 / 140**

## **Discussion**

**WG2 / 137**

## **Discussion**

**Session 2 / 202**

## **Discussion: (re)forming a nuSTORM collaboration**

**WG4 / 120**

## **Distinguishing muon LFV effective couplings using $\mu \rightarrow e e \gamma$**

**Corresponding Author(s):** joe@phy.saitama-u.ac.jp

**Posters & welcome reception / 45**

## **Distinguishing muon LFV effective couplings using $\mu \rightarrow e e \gamma$**

Joe Sato<sup>1</sup>

<sup>1</sup> *Saitama University*

**Corresponding Author(s):** joe@phy.saitama-u.ac.jp

We discuss how to discriminate muon LFV couplings one from the other using the mode  $\mu \rightarrow e e \gamma$ .

**WG1 Neutrino oscillation / 173**

## **Double Chooz**

**WG3 Parallel Session 5 / 158**

## **EMuS at CSNS Updated Studies**

**Corresponding Author(s):** vassilopoulos@ihep.ac.cn

**WG3 Parallel Session 1 / 145**

## **ESSnuSB Target and Horn Studies and Future Development**

**Corresponding Author(s):** marcos.dracos@in2p3.fr

**Plenary V / 24**

## **Electron Scattering**

**Corresponding Author(s):** vishvas@vt.edu

**WG2 / 131**

## **Electron scattering data and neutrinos**

**Corresponding Author(s):** weinstein@odu.edu

**WG2 / 78**

## **Electron vs Muon Neutrinos**

**Corresponding Author(s):** artank@vt.edu

**WG5 / 93**

## **Exploring the Potential of Short-Baseline Physics at Fermilab**

**Corresponding Author(s):** pedrosimpas@gmail.com

**Plenary VII / 18**

## **FNAL SBL Program Status**

**Corresponding Author(s):** afava@fnal.gov

**WG1+WG2 / 76**

## FNAL SBN Status

**Corresponding Author(s):** jaz8600@fnal.gov

**Posters & welcome reception / 50**

### **Fabrication of A High-efficiency Cosmic Ray Veto Detector for the Mu2e Experiment**

**Author(s):** Ben Barton<sup>1</sup> ; Danny Mills<sup>1</sup>

**Co-author(s):** Craig Group<sup>2</sup>

<sup>1</sup> *Virginia*

<sup>2</sup> *University of Virginia*

**Corresponding Author(s):** dm8bz@virginia.edu, bb6yx@virginia.edu, group@virginia.edu

The Mu2e experiment at Fermilab will search for the charged-lepton-flavor-violating process of coherent muon-to-electron conversion in the presence of a nucleus with a sensitivity four orders of magnitude beyond the current strongest limits. The goal of single-event sensitivity requires that all backgrounds must sum to significantly less than one event. One potential background is due to cosmic-ray muons producing an electron with signal characteristics within the Mu2e apparatus. The cosmic-ray-veto system of the Mu2e experiment is tasked with vetoing such cosmic-ray-induced backgrounds with high efficiency while inducing low dead time and while operating in the high-intensity environment of the Mu2e experiment. The UVA HEP group has been leading the effort to design and prototype the CRV and has recently started the fabrication this detector on site. Highlights of this effort including production detector performance will be presented.

**WG1 Neutrino oscillation / 181**

### **Final results from the OPERA experiment in the CNGS neutrino beam**

**Corresponding Author(s):** tenti@bo.infn.it

**WG1 / 172**

### **Final results from the OPERA experiment in the CNGS neutrino beam**

**Corresponding Author(s):** tenti@bo.infn.it

**Posters & welcome reception / 43**

### **Final results from the OPERA experiment in the CNGS neutrino beam**

Matteo Tenti<sup>1</sup>

<sup>1</sup> INFN**Corresponding Author(s):** matteo.tenti@cnaf.infn.it

The OPERA experiment at the Gran Sasso Laboratory was designed to study  $\nu_{\mu} \rightarrow \nu_{\tau}$  oscillations in appearance mode in the CERN-to-Gran Sasso neutrino beam. We report the final analysis of the full data sample based on looser selection criteria than in previous analyses and multivariate approach. Oscillation parameters have been determined with a reduced statistical uncertainty, and the discovery of tau neutrino appearance is confirmed with an improved significance level. Moreover, the search for electron neutrino events has been extended to the full dataset exploiting an improved method for the electron neutrino energy estimation. New limits have been set in the 3+1 neutrino model.

WG1+WG5 / 193

## First Results from the PROSPECT Short Baseline Reactor Experiment

**Corresponding Author(s):** nbowden@llnl.gov

WG5 / 99

## Future DUNE constraints on EFT

**Corresponding Author(s):** ggrilli@if.usp.br

WG3 Parallel Session 1 / 143

## Future Upgrade of J-PARC Target and Beam Window

**Corresponding Author(s):** chris.densham@stfc.ac.uk

Posters &amp; welcome reception / 64

## Future prospects for CAPTAIN experiment

Jorge Chaves<sup>1</sup><sup>1</sup> University of Pennsylvania**Corresponding Author(s):** jchaves@sas.upenn.edu

The CAPTAIN (Cryogenic Apparatus for Precision tests of Argon Interactions with Neutrinos) experiment is a five-ton liquid argon time projection chamber (LArTPC) at Los Alamos National Laboratory. CAPTAIN is designed to make measurements of liquid argon interactions relevant to neutrino physics in particular for the proposed Deep Underground Neutrino Experiment (DUNE). A prototype detector called Mini-CAPTAIN, with 400 kg of liquid argon, collected data at a neutron beam at LANL in the summer of 2017. We present plans for the future of the CAPTAIN experiment to take data at other neutrino sources and measure low-energy neutrino interactions on argon.

**WG2 / 136**

## **GENIE Physics Tuning**

**Corresponding Author(s):** ljiang@pitt.edu

**Plenary X / 34**

## **Global Neutrino Oscillation Fits**

**Corresponding Author(s):** gsk2112@columbia.edu

**WG1 Neutrino oscillation / 184**

## **Global analysis of neutrino oscillation experiments**

**Corresponding Author(s):** chternes@ific.uv.es

**WG3 Parallel Session 4 / 154**

## **Integrable Optics Test Accelerator**

**Corresponding Author(s):** bfreemire@niu.edu

**WG3 Parallel Session 7 / 156**

## **IsoDAR**

**Corresponding Author(s):** smolsky@mit.edu

**WG2 / 73**

## **LArIAT Cross Section Results**

**Corresponding Author(s):** mtzanov@fnal.gov

**Plenary VII / 17**

## **LHC Results on heavy Neutrino Searches**

**Corresponding Author(s):** ukyang@snu.ac.kr

**WG1+WG5 / 191**

## **Latest Results from MINOS+ on Sterile Neutrinos search**

**Corresponding Author(s):** toddjb@mail.uc.edu

**WG1 Neutrino oscillation / 175**

## **Latest Results from the Daya Bay Reactor Neutrino Experiment**

**Corresponding Author(s):** goowenq@gmail.com

**WG2 / 70**

## **Lattice QCD and neutrinos**

**Corresponding Author(s):** ask@fnal.gov

**WG4 - CLFV at Colliders / 115**

## **Lepton Flavour Universal at LHCb**

**Corresponding Author(s):** francesca.dordei@cern.ch

**WG5 / 95**

## **Lepton-Number-Charged Scalars and Neutrino Beamstrahlung**

**Corresponding Author(s):** kevinkelly2019@u.northwestern.edu

**WG5 / 90**

## **Light scalar dark matter at neutrino oscillation experiments**

**Corresponding Author(s):** whisnant@iastate.edu

**Plenary VIII / 23**

## **Low Emittance Muon Accelerator**

**Corresponding Author(s):** manuela.boscolo@lnf.infn.it

**WG3 Parallel Session 4 / 162**

## **Low Emittance Muon Beams**

**Corresponding Author(s):** manuela.boscolo@lnf.infn.it

**WG2 / 133**

## **Low energy neutrino interactions**

**Corresponding Author(s):** sch52@phy.duke.edu

**Plenary VIII / 22**

## **MICE Results**

**Corresponding Author(s):** paul.soler@glasgow.ac.uk

**WG1+WG2 / 74**

## **MINERvA Cross Section Results**

**Corresponding Author(s):** xianguo.lu@physics.ox.ac.uk

**WG3 Parallel Session 2 / 148**

## **Measurement of Phase-Space Density Evolution in MICE**

**Corresponding Author(s):** torun@iit.edu

**WG5 / 194**

## **Measuring the Leptonic Dirac CP Phase with Muon Decay at Rest**

**Corresponding Author(s):** gesf02@gmail.com

**WG1+WG2 / 75**

## **MicroBooNE Cross Section Results**

**Corresponding Author(s):** libojiang8584@gmail.com



**WG1 Neutrino oscillation / 185**

## **MicroBoONE Search for Low-Energy Excess Using Deep Learning Algorithms**

**Corresponding Author(s):** lyates@mit.edu**Posters & welcome reception / 63**

## **Mini-CAPTAIN measurements in the LANSCE WNR Neutron Beam**

Clark McGrew<sup>1</sup><sup>1</sup> *Stony Brook University***Corresponding Author(s):** clark.mcgrew@stonybrook.edu

All neutrino experiments face the problem of reconstructing the incoming neutrino energy using the visible interaction products. Unfortunately, the initial neutrino interaction is not well understood, not all of the interaction products are visible, and the secondary interactions may not be well understood. In preparation the analysis of neutrino oscillation data collected using liquid argon time projection chambers, the CAPTAIN collaboration is addressing this problem with a measurement of the cross section of neutrons impinging on an argon target. Using the WNR neutron facility, which produces a well known flux of neutrons up to a kinetic energy of 800 MeV, the total cross section will be measured for neutron kinetic energies above approximately 50 MeV, and partial cross sections will be measured for  $n + \text{Ar} \rightarrow p + X$  and  $n + \text{Ar} \rightarrow \pi^\pm + X$ . Data for this measurement was collected during the Summer of 2017 using a 400 kg fiducial Liquid Argon TPC that was instrumented with a photon-detection system (PDS). The interaction by interaction neutron energy is determined using time of flight as determined by the PDS while the ionization yield is measured in the TPC.

**WG5 / 92**

## **Model independent non-unitarity**

**Corresponding Author(s):** hisakazu.minakata@gmail.com**Posters & welcome reception / 44**

## **Modeling neutrino-nucleus interactions in the few-GeV region**

**Author(s):** Kajetan Niewczas<sup>1</sup>**Co-author(s):** Alexis Nikolakopoulos<sup>1</sup>; James Nys<sup>1</sup>; Natalie Jachowicz<sup>1</sup>; Nils Van Dessel<sup>1</sup>; Raúl González-Jiménez<sup>2</sup>; Vishvas Pandey<sup>3</sup><sup>1</sup> *Ghent University*<sup>2</sup> *Complutense University of Madrid*<sup>3</sup> *Virginia Tech*

**Corresponding Author(s):** kajetan.niewczas@ugent.be, vishvas@vt.edu, raugonjim@gmail.com, natalie.jachowicz@ugent.be, nils.vandessel@ugent.be, jannes.nys@ugent.be, alexis.nikolakopoulos@ugent.be

A good understanding of neutrino-nucleus scattering mechanisms is essential to reduce the systematic errors in neutrino oscillation experiments. Recent interest of the Ghent group focus on providing a consistent description of this process in the intermediate energy region. We describe the low energy response with collective nuclear excitations and the quasielastic peak using a Hartree-Fock-CRPA (continuum random phase approximation) model that takes into account nuclear long-range correlations as well as hadronic final-state interactions. The two-body current mechanisms, which are especially important in the region between the quasielastic and the delta-resonance peak, are included through short-range correlations and meson-exchange currents, treated within the same mean-field based model. Our description of intermediate-energy neutrino-nucleus scattering is completed by modeling neutrino-induced pion production. For that, we consider the dominant contribution from the decay of the delta resonance as well as other terms required by chiral symmetry, working in a fully relativistic formalism with a refined treatment of nuclear effects.

**WG4 - CLFV 2 / 108**

## **Mu2e II - A Proposed Evolution of the Mu2e Experiment**

**Corresponding Author(s):** group@virginia.edu

**Posters & welcome reception / 59**

## **Muonic X-ray measurements with radioactive elements**

Frederik Wauters<sup>1</sup> ; Alexander Skawran<sup>2</sup>

<sup>1</sup> *Johannes Gutenberg Universitaet Mainz*

<sup>2</sup> *ETH*

**Corresponding Author(s):** fwauters@uni-mainz.de, alexander.skawran@psi.ch

Muonic X-rays are an excellent tool to measure the nuclear charge radius, the radii of almost all stable nuclei have been measured with this method. More challenging are radioactive nuclei due to the amount of material needed to stop negative muons produces at accelerator facilities. At the Paul Scherrer Institute we have developed a novel method, stopping the muons in a gaseous hydrogen-deuterium mixture. Initially muonic hydrogen is formed, then the muon transfers to deuterium. Due to a minimum in the scattering cross section, this muonic deuterium quickly reaches the target chambers walls, where the muons transfer a higher Z element. A layer as thin as a few nanometers of the element of interest is sufficient to produce the muonic atoms with a efficiency of O(10%). We aim to measure the muonic X-rays in <sup>226</sup>Ra. The charge radius derived from this data will serve as a crucial input for an upcoming atomic parity violation experiment with a single trapped radium atom.

**WG2 / 134**

## **NOvA Cross Section Model / Oscillation Needs**

**Corresponding Author(s):** jeremy.wolcott@tufts.edu

WG2 / 71

## NOvA Cross Section Results

Corresponding Author(s): matthew.judah@colostate.edu

WG2 / 139

## NUISANCE for neutrino cross section fits

Corresponding Author(s): picker24@msu.edu

Plenary X / 35

## Neutrino Dark Matter Connections

Corresponding Author(s): krnjaicg@fnal.gov

WG5 / 89

## Neutrino Oscillations in Dark Backgrounds

Corresponding Author(s): vecchi.alsz@gmail.com

Posters &amp; welcome reception / 206

## Neutrino Physics with Deep Learning on NOvA

Fernanda Psihas<sup>None</sup>

Corresponding Author(s): psahas@fnal.gov

The NOvA experiment has made both  $\nu_\mu$  disappearance and  $\nu_e$  appearance measurements in Fermilab's NuMI beam, and is working on cross section measurements using near detector data. At the core of NOvA's measurements is the use of deep learning algorithms for identification and reconstruction of the neutrino flavor and energy.

Presented here is the extension of our deep learning efforts for identification of neutrino signal events, final state identification, single particle tagging, and reconstruction using instance segmentation techniques. I will describe the new implementations of modified Convolutional Neural Networks for anti-neutrino events which yield a 14% improvement in efficiency. I will also show the performance of our single particle ID network, data driven performance tests, standard candle measurements, and advances for reconstruction.

Posters &amp; welcome reception / 161

## Neutrino Physics with Deep Learning on NOvA

Fernanda Psihas<sup>None</sup>

**Corresponding Author(s):** psihas@fnal.gov

The NOvA experiment has made both  $\nu_\mu$  disappearance and  $\nu_e$  appearance measurements in Fermilab's NuMI beam, and is working on cross section measurements using near detector data. At the core of NOvA's measurements is the use of deep learning algorithms for identification and reconstruction of the neutrino flavor and energy.

Presented here is the extension of our deep learning efforts for identification of neutrino signal events, final state identification, single particle tagging, and reconstruction using instance segmentation techniques. I will describe the new implementations of modified Convolutional Neural Networks for anti-neutrino events which yield a 14% improvement in efficiency. I will also show the performance of our single particle ID network, data driven performance tests, standard candle measurements, and advances for reconstruction.

WG5 / 85

## Neutrino flavor transformation in supernova as a probe for non-standard neutrino-scalar interactions

**Corresponding Author(s):** jim\_kneller@ncsu.edu

WG1 Neutrino oscillation / 180

## Neutrino physics with KM3NeT/ORCA

**Corresponding Author(s):** zaborov@cppm.in2p3.fr

WG1 / 171

## Neutrino physics with KM3NeT/ORCA

**Corresponding Author(s):** zaborov@cppm.in2p3.fr

WG5 / 101

## Neutrino trident at DUNE

**Corresponding Author(s):** matheus.hostert@durham.ac.uk

Posters & welcome reception / 53

## Neutrino trident production at near detectors

Matheus Hostert<sup>1</sup>

<sup>1</sup> *IPPP, Durham University*

**Corresponding Author(s):** matheus.hostert@durham.ac.uk

The large statistics expected at the near detectors of neutrino oscillation experiments opens up the possibility to search for rare neutrino interactions. One example is neutrino trident production, the scattering of a neutrino by the Coulomb field of a nucleus producing a pair of charged leptons. In this talk, I will revisit the calculation of the trident scattering rate, addressing certain inconsistencies in the literature and presenting revised predictions for the total and differential event rates for relevant experiments. I will then argue that backgrounds can be kept under control and that certain channels could be seen for the first time at these facilities. Finally, I will dedicate some time to discuss what kind of new physics one can look for in these processes.

**WG1 Neutrino oscillation / 174**

## **New Results from RENO**

**Corresponding Author(s):** jsjang@gist.ac.kr

**WG2 / 67**

## **Nuclear Theory, Data and Event Generators**

**Corresponding Author(s):** jvanorde@odu.edu

**WG2 / 138**

## **Nucleon axial form-factor from a Bayesian neural network analysis of scattering data**

**Corresponding Author(s):** edusaul@ific.uv.es

**WG2 / 77**

## **Open Issues in Nuclear SIS and DIS Scattering**

**Corresponding Author(s):** morfin@fnal.gov

**Plenary I / 2**

## **Overview of Neutrino Physics**

**Corresponding Author(s):** ichikawa@scphys.kyoto-u.ac.jp

**Posters & welcome reception / 49****Performance of Scintillation Counters as Measured at the Fermilab Test Beam Facility for the Mu2e Cosmic Ray Veto System****Author(s):** Ningshun Chen<sup>1</sup>**Co-author(s):** Craig Group<sup>2</sup><sup>1</sup> *Virginia*<sup>2</sup> *University of Virginia***Corresponding Author(s):** nc2bx@virginia.edu, group@virginia.edu

Photoelectron yields of extruded scintillation counters with titanium dioxide coating and embedded wavelength shifting fibers read out by silicon photomultipliers have been measured at the Fermilab Test Beam Facility using 120 GeV protons. The yields were measured as a function of transverse, longitudinal, and angular positions for a variety of scintillator compositions, reflective coating mixtures, and fiber diameters. Timing performance was also studied. These studies were carried out by the Cosmic Ray Veto Group of the Mu2e collaboration as part of their R&D program.

**Posters & welcome reception / 51****Performance of Wavelength-Shifting Fibers for the Mu2e Cosmic Ray Veto Detector****Author(s):** Peter Farris<sup>1</sup>**Co-author(s):** Craig Group<sup>2</sup><sup>1</sup> *Virginia*<sup>2</sup> *University of Virginia***Corresponding Author(s):** pjf4qr@virginia.edu, group@virginia.edu

The Mu2e experiment will search for a neutrino-less muon-to-electron conversion process with almost four orders of magnitude of sensitivity improvement relative to the current best limit. One important background is caused by cosmic ray muons, and particles produced by their decay or interactions, mimicking the conversion electron signature. In order to reach the design sensitivity, Mu2e needs to obtain a cosmic ray veto (CRV) efficiency of 99.99%. The CRV system consists of four layers of plastic scintillating counters read out by silicon photo-multipliers (SiPM) through wavelength shifting fibers. The CRV counters must produce sufficient photo statistics in order to achieve the required veto efficiency. We study the light properties of several wavelength shifting fiber sizes in order to optimize the total light yield for the CRV system. The measurements are performed using a scanner designed to ensure fiber quality for the CRV. Results from prototype and production fiber studies will be presented.

**WG1 Neutrino oscillation / 187****Physics potential of Hyper-Kamiokande for neutrino oscillation measurements****Corresponding Author(s):** tetsuro.sekiguchi@kek.jp

**WG1 Neutrino oscillation / 188**

## **Physics potential of the ESS $\nu$ SB facility**

**Corresponding Author(s):** salvador.rosauro@uam.es

**WG4 - Muonic atoms/proton radius / 124**

## **Precision spectroscopy of exotic atoms involving muon**

**Corresponding Author(s):** sohtaro.kanda@riken.jp

**WG1 Neutrino oscillation / 176**

## **Probing Neutrino Mass Ordering and Solar neutrinos with JUNO detector**

**Corresponding Author(s):** xuefeng.ding@gssi.it

**WG5 / 81**

## **Probing neutrino coupling to a light scalar with coherent neutrino scattering**

**Corresponding Author(s):** xunjie.xu@gmail.com

**WG5 / 88**

## **Probing secret interactions of eV-scale sterile neutrinos with the diffuse supernova neutrino background**

**Corresponding Author(s):** mary-hall-reno@uiowa.edu

**Plenary IX / 31**

## **Pulsed Muon Beam Physics**

**Corresponding Author(s):** m.lancaster@ucl.ac.uk

**Posters & welcome reception / 207**

## **Reactor Antineutrino Detection Using CHANDLER: A New Portable Neutrino Detector** Tulasi Subedi **Abstract CHANDLER**

Tulasi Subedi<sup>None</sup>

**Corresponding Author(s):** tpsubedi@vt.edu

CHANDLER is a neutrino detection technology to detect reactor antineutrino. It detects the end products (positron and neutron) from inverse beta decay (IBD) reaction, to tag an event. This technology can be used for nuclear non-proliferation and a sterile neutrino search.

**Posters & welcome reception / 61**

## **Recent Cross Section Results from T2K**

Clarence Wret<sup>1</sup>

<sup>1</sup> *University of Rochester*

**Corresponding Author(s):** c.wret14@imperial.ac.uk

Measurements of the PMNS oscillation parameters by the T2K experiment are improving our understanding of neutrino mixing. Using the two multi-purpose near detectors, ND280 and INGRID, T2K also extensively measures neutrino-nucleus interactions in the low GeV region. With multiple targets and on/off-axis detector placement, the near detectors investigate target dependence and the effects of different neutrino fluxes. This talk introduces T2K and its cross section measurements, with emphasis on recent and upcoming results.

**WG3 Parallel Session 1 / 142**

## **Recent Progress on Radiation Damage Studies at RaDIATE**

**Corresponding Author(s):** taku.ishida@kek.jp

**WG3 Parallel Session 2 / 146**

## **Recent Results from MICE on Multiple Coulomb Scattering and Energy Loss**

**Corresponding Author(s):** john.nugent@glasgow.ac.uk

**Plenary V / 25**

## **Recent Results from MINERVA**

**Corresponding Author(s):** mateuscarneiro@gmail.com



WG3 Parallel Session 2 / 147

## Recent Results from the Study of Emittance Evolution in MICE

**Corresponding Author(s):** christopher.hunt08@imperial.ac.uk

Plenary VI / 27

## Recent Results from the T2K Near Detector

**Corresponding Author(s):** xianguo.lu@physics.ox.ac.uk

Posters &amp; welcome reception / 66

## Recent progress on radiation damage studies at RaDIATE

**Author(s):** Taku Ishida<sup>1</sup>

**Co-author(s):** Patrick G. Hurh<sup>2</sup>

<sup>1</sup> *J-PARC/KEK*

<sup>2</sup> *FNAL*

**Corresponding Author(s):** taku.ishida@kek.jp, hurh@fnal.gov

In the recent past, major accelerator facilities have been limited in beam power not by their accelerators but by target and/or window survivability. With present plans to upgrade accelerator facilities at FNAL and J-PARC to higher beam powers (1.2+ MW) in the next decade, timely R&D of robust high power targets and beam windows is needed to fully realize the physics benefits of the higher beam power. An international team of researchers, under the aegis of the Radiation Damage In Accelerator Target Environments (RaDIATE) Collaboration, fabricated test specimens which were irradiated by 181 MeV protons in the Brookhaven Linac Isotope Producer (BLIP) facility at BNL, starting in spring of 2017. Test specimens, including candidate materials for various beam intercepting device applications, were provided by participating facilities. Post-irradiation examination (PIE) is being conducted at participating RaDIATE institutions with appropriate “hot-cell” facilities. The work includes efforts to provide BLIP irradiated samples to in-beam thermal shock test at CERN’s HiRadMat beam-line facility. Thermal shock testing in beam allows observation of how the radiation damaged property data affects material behavior when exposed to actual beam loading conditions. The HiRadMat beam-line experiment proposal was accepted by the HiRadMat Scientific Board and is currently scheduled to run in October, 2018. In this talk up-to-date status of the experiments, PIEs, and prospect for the works conducted by RaDIATE collaboration will be over-viewed.

Posters &amp; welcome reception / 56

## Recent results of neutrino interactions from the T2K Near Detector

Xianguo Lu<sup>None</sup>

**Corresponding Author(s):** xianguo.lu@physics.ox.ac.uk

Neutrino-nucleus cross-section measurements in the GeV regime are crucial for future accelerator-based precision neutrino oscillation measurements. The T2K Near Detector has provided important results for the study of nuclear effects in neutrino-nucleus interactions and therefore stringent constraints on model development. In this talk, I will present our recent cross-section measurements, highlight on-going progress and discuss future possible developments in T2K.

WG2 / 80

## **Removal and Binding Energy in Lepton-Nucleus Scattering**

Corresponding Author(s): bodek@pas.rochester.edu

Plenary III / 12

## **Results and Prospects from NOvA**

Corresponding Author(s): bianjm@physics.umn.edu

Plenary IV / 13

## **Results and Prospects from T2K**

Corresponding Author(s): s.r.dennis@liverpool.ac.uk

Plenary IV / 14

## **Results from IceCube**

Corresponding Author(s): caad@mit.edu

Plenary V / 26

## **Review of Tension in Data/Models of Neutrino Cross Sections**

Corresponding Author(s): mahn@pa.msu.edu

Plenary IX / 33

## **SBL Reactor Experiments**

Corresponding Author(s): chao@bnl.gov

**Posters & welcome reception / 54****Search for  $K^+ \rightarrow \pi^+ \nu \nu$  at CERN**Stoyan Trilov<sup>None</sup>

Corresponding Author(s): stoyan.trilov@bristol.ac.uk

The decay  $K^+ \rightarrow \pi^+ \nu \nu$ , with a very precisely predicted branching ratio of less than  $10^{-10}$ , is one of the best candidates to reveal indirect effects of new physics at the highest mass scales. The NA62 experiment at CERN SPS is designed to measure the branching ratio of the  $K^+ \rightarrow \pi^+ \nu \nu$  with a decay-in-flight technique, novel for this channel. NA62 took data in 2016, 2017 and another year run is scheduled in 2018. Statistics collected in 2016 allows NA62 to reach the Standard Model sensitivity for  $K^+ \rightarrow \pi^+ \nu \nu$ , entering the domain of  $10^{-10}$  single event sensitivity and showing the proof of principle of the experiment. The analysis data is reviewed and the preliminary result from the 2016 data set presented.

**WG4 - CLFV at Colliders / 113****Search for lepton flavour violation with the ATLAS detector**

Corresponding Author(s): ws.chan@cern.ch

**WG4 / 119****Searches for Electric Dipole Moments (EDM) at a Storage Ring with JEDI**

Corresponding Author(s): m.zurek@fz-juelich.de

**WG4 / 121****Searches for heavy neutral lepton production and lepton flavour violation in kaon decays at the NA62 experiment**

Corresponding Author(s): stoyan.trilov@bristol.ac.uk

**Plenary VII / 141****Shedding light on low energy excess anomaly with MiniBooNE and MicroBooNE**

Corresponding Author(s): zarko@fnal.gov

**Session 2 / 201**

## **Simulation studies of a detector for nuSTORM**

**Corresponding Author(s):** p.hallsjo.1@research.gla.ac.uk

WG5 / 97

## **Single Ion Barium Tagging for Neutrino-less Double-Beta Decay Searches. A multi-disciplinary technique in development for the NEXT experiment**

**Corresponding Author(s):** psihas@fnal.gov

Session 1 / 196

## **Sketch of physics case in preparation for the EU Strategy Update**

**Corresponding Author(s):** k.long@imperial.ac.uk

WG5 / 100

## **Solar nu at DUNE and BSM**

**Corresponding Author(s):** shirleyl@slac.stanford.edu

WG4 - g-2, precision physics with muons / 111

## **Standard Model prediction for the muon g-2**

**Corresponding Author(s):** dnomura@yukawa.kyoto-u.ac.jp

WG5 / 86

## **Standard and non-standard neutrino physics at reactor experiments**

**Corresponding Author(s):** dvanegas@vt.edu

Plenary III / 11

## **Status and Future of high-power Neutrino Target Stations**

**Corresponding Author(s):** hylen@fnal.gov

Plenary III / 10

## **Status and Future of high-power Proton Drivers**

**Corresponding Author(s):** scousine@ornl.gov

WG3 Parallel Session 1 / 144

## **Status and Physics Potential of MOMENT Study**

**Corresponding Author(s):** tangjian5@mail.sysu.edu.cn

Plenary IV / 15

## **Status and Physics of JUNO**

**Corresponding Author(s):** xuefeng.ding@gssi.it

WG4 - CLFV at Colliders / 114

## **Status of Charged Lepton Flavor Violation searches at CMS and future prospects**

**Corresponding Author(s):** diego.beghin@ulb.ac.be

Plenary VIII / 20

## **Status of DUNE**

**Corresponding Author(s):** sgollapi@utk.edu

Plenary VIII / 21

## **Status of ESSnuB**

**Corresponding Author(s):** marcos.dracos@in2p3.fr

WG3 Parallel Session 5 / 159

## **Status of NA61/SHINE Measurements for Neutrino Experiments**

**Corresponding Author(s):** dathula@gmail.com

WG1 Neutrino oscillation / 189

## **Status of ProtoDUNE Experiments at CERN**

**Corresponding Author(s):** jiowang@ucdavis.edu

Plenary VII / 19

## **Status of T2HK**

**Corresponding Author(s):** alain.blondel@cern.ch

Plenary I / 3

## **Status of accelerator-based Neutrino Physics**

**Corresponding Author(s):** plvahle@wm.edu

Session 1 / 199

## **Status of consideration of implementation of nuSTORM at CERN**

**Corresponding Author(s):** mike.lamont@cern.ch

WG4 -CLFV I / 106

## **Status of the COMET Experiment**

**Corresponding Author(s):** moritsu@post.kek.jp

WG4 -CLFV I / 105

## **Status of the DeeMe Experiment**

**Corresponding Author(s):** d-nagao@kuno-g.phys.sci.osaka-u.ac.jp

**WG3 Parallel Session 5 / 160**

## **Status of the ENUBET**

**Corresponding Author(s):** brunetti@pd.infn.it, giulia.brunetti@cern.ch

**WG3 and WG4 - Muon Beam Facilities / 126**

## **Status of the Facility/Accelerator/Beam-line for Muon Programs at J-PARC**

**Corresponding Author(s):** hajime.nishiguchi@kek.jp

**WG5 / 94**

## **Status of the ISODAR project**

**Corresponding Author(s):** saxani@mit.edu

**WG4 -CLFV I / 103**

## **Status of the MEG II Experiment**

**Corresponding Author(s):** angela.papa@psi.ch

**WG1+WG5 / 190**

## **Sterile Neutrinos search via NC dis at NOvA**

**WG1+WG5 / 192**

## **Sterile neutrino searches with the ICARUS detector**

**Corresponding Author(s):** yuntse@slac.stanford.edu

**WG5 / 96**

## **Sterile neutrinos searches**

**Corresponding Author(s):** josu.hernandez@sissa.it

WG4 - CLFV 2 / 109

## Studies of PRISM/PRIME - the next generation muon to electron conversion experiment

Corresponding Author(s): j.pasternak@imperial.ac.uk

Posters &amp; welcome reception / 52

## Studies of the Aging Properties of the Mu2e Cosmic Ray Veto System

Author(s): Yuri Oksuzian<sup>1</sup>

Co-author(s): Craig Group<sup>2</sup>

<sup>1</sup> Virginia

<sup>2</sup> University of Virginia

Corresponding Author(s): oksuzian@fnal.gov, group@virginia.edu

The Muon-to-Electron Conversion experiment (Mu2e) operates at extremely high sensitivities, requiring a means of reducing experimental background. The Cosmic Ray Veto system (CRV) is a particle detector that will surround the Mu2e apparatus to veto penetrating particles that present background. The CRV must have a detection efficiency of 99.99% throughout the expected three year lifetime of the Mu2e experiment. The CRV is comprised of extruded polystyrene scintillating strips and fiber which degrade over time, decreasing the efficiency of the CRV. Using a standard accelerated aging technique, several scintillator and fiber samples were heated to increase their rate of degradation. The results of these studies and the impact of aging on the CRV will be presented.

Posters &amp; welcome reception / 65

## Studying Neutral Current Elastic Scattering and the Strange Axial Form Factor in MicroBooNE

Katherine Woodruff<sup>1</sup>

<sup>1</sup> New Mexico State University

Corresponding Author(s): kwoodruf@nmsu.edu

One of the least constrained contributions to the neutral current (NC) elastic neutrino-proton cross section is the strange axial form factor, which represents the strange quark spin contribution to the spin of the proton. Knowledge of this form factor is important for many areas of physics including sterile neutrino searches, spin-dependent dark matter searches, and supernova explosion mechanisms. The strange axial form factor can be determined by studying NC elastic scattering events in the MicroBooNE detector at low negative four-momentum transfer squared ( $Q^2$ ). MicroBooNE's unique ability to detect low-energy protons is expected to allow the measurement of these events with a  $Q^2$  as low as 0.10 GeV<sup>2</sup>. We present a selection of neutral current elastic events in a subset of MicroBooNE neutrino data, as well as our plan to extract the strange axial form factor from this selection in the full data set.



WG2 / 135

## **T2K Cross Section Model / Oscillation Needs**

Corresponding Author(s): c.wret14@imperial.ac.uk

WG2 / 72

## **T2K Cross Section Results**

Corresponding Author(s): c.wret14@imperial.ac.uk

Plenary VI / 28

## **T2K Near Detector Upgrades and Plans for T2HK**

Corresponding Author(s): thorsten.lux@ifae.es

WG4 / 118

## **Test on lepton flavor universality at BESIII**

WG3 Parallel Session 3 / 149

## **The LBNF Beamline**

Corresponding Author(s): mbishai@bnl.gov

WG4 - Muonic atoms/proton radius / 122

## **The MUon Scattering Experiment (MUSE) at the Paul Scherrer Institute**

Corresponding Author(s): strauch@sc.edu

WG4 - CLFV 2 / 107

## **The Mu2e Experiment at Fermilab**

Corresponding Author(s): sb8es@virginia.edu

**Posters & welcome reception / 47****The Mu2e Experiment at Fermilab****Author(s):** Steve Bui<sup>1</sup>**Co-author(s):** Craig Group<sup>2</sup><sup>1</sup> *Virginia*<sup>2</sup> *University of Virginia***Corresponding Author(s):** sb8es@virginia.edu, group@virginia.edu

The Muon-to-Electron-Conversion (Mu2e) Experiment is a high-precision, intensity-frontier experiment being developed at Fermilab which will search for coherent, neutrino-less muon to electron conversion in the presence of an atomic nucleus. Such a process would exhibit charged lepton flavor violation (CLFV), which has not yet been observed. Continuing the search for CLFV, Mu2e will improve the sensitivity by four orders of magnitude over the present limits. In the search for beyond the standard model (BSM) physics, Mu2e is uniquely sensitive to a wide range of models by indirectly probing mass scales up to the energy scale of  $10^4$  TeV. While muon-to-electron-conversion is permissible in the standard model through neutrino oscillations, the rate is extremely low at about one event in  $10^{52}$ . By design, the background for the experiment will be well-understood and kept at a sub-event level, which will mean the observation of muon-to-electron conversion is a direct confirmation of BSM physics. The physics motivation, the design, and the current status of the experiment will be presented.

**Posters & welcome reception / 205****The NOvA Test Beam Program**Andrew Sutton<sup>None</sup>**Corresponding Author(s):** ats2mk@virginia.edu

NOvA is a long-baseline off-axis beam neutrino experiment. By measuring  $\nu_\mu$  disappearance and  $\nu_e$  appearance at the 14 kiloton NOvA Far Detector, the experiment is addressing outstanding questions in neutrino physics, including the neutrino mass hierarchy and existence of leptonic CP violation. The NOvA Test Beam program, under deployment at the Fermilab Test Beam Facility, will use a scaled-down NOvA detector to sample beams of tagged electrons, muons, pions, and protons in the momentum range of 0.3 to 2 GeV/c. It will further the NOvA physics reach by precisely measuring the detector's muon energy scale and electromagnetic and hadronic response, and provide real data for detailed studies of particle identification techniques. Ongoing efforts on beamline instrumentation, data acquisition, simulation, momentum reconstruction and particle identification are presented. Implications for the neutrino oscillation measurements are discussed.

**Posters & welcome reception / 62****The Role of Cross Sections in the Oscillation Analysis: The T2K Experience**Clarence Wret<sup>1</sup><sup>1</sup> *University of Rochester***Corresponding Author(s):** c.wret14@imperial.ac.uk

The T2K experiment measures long baseline neutrino oscillations with neutrinos in the 0.1-1.5 GeV energy range. Thanks to excellent beam performance T2K is rapidly gathering statistics, increasing the relative importance of the parameterisation of systematics. Neutrino-nucleus interactions are large contributors to the error budget at T2K, affecting crucial components such as neutrino energy estimation and event selection. This talk gives an overview of T2K's treatment of interaction systematics, the constraints that are placed upon them, and their impact on oscillation analyses.

**WG1 Neutrino oscillation / 177**

## **The design and research progresses of the Central Detector in JUNO**

**Corresponding Author(s):** henyk@ihep.ac.cn

**WG3 Parallel Session 7 / 155**

## **The g-2 Beamline**

**Corresponding Author(s):** nfroemm@fnal.gov

**WG4 - g-2, precision physics with muons / 110**

## **The g-2 experiment at Fermilab**

**Corresponding Author(s):** kaspar@uw.edu

**WG4 -CLFV I / 104**

## **The search for Lepton-Flavour Violation with the Mu3e Experiment**

**Corresponding Author(s):** fwauters@uni-mainz.de

**Posters & welcome reception / 55**

## **The search for neutral currents in muonic X-rays**

Frederik Wauters<sup>1</sup>

<sup>1</sup> *Johannes Gutenberg Universitaet Mainz*

**Corresponding Author(s):** fwauters@uni-mainz.de

### Muonic X-ray measurements at the Paul Scherrer Institute

Negative muons at rest quickly get captured by nearby atoms in highly excited atomic states. These muonic atoms subsequently de-excite via radiative and Auger transitions until the muon ends up in the 1s orbital. At the lower orbits, there is substantial overlap between the muon wave function and the nucleus, making this system an excellent laboratory to study the interaction between the muon and atomic nucleus.

MuX is a renewed effort at the Paul Scherrer to measure muonic X-rays in medium- and high-Z nuclei, fully exploiting the coverage and multiplicity of a germanium detector array and the high yield of negative muons available. The physics program focuses on atomic parity violation (APV). A measurement of the charge radius of  $^{226}\text{Ra}$ , derived from the 2s-1s transition energy, will serve as crucial input for an upcoming APV experiment with a single Ra ion. A second measurement program is exploring the possibility of measuring APV directly in muonic atoms. In the Standard Model, APV arises from the mixing of the opposite parity 2p and 2s atomic states, leading to parity violation in the 2s-1s transition. We focus on  $Z=30$  nuclei, where a measurable branching ratio of the single photon 2s-1s transition is expected. The high granularity of a large solid angle germanium detector array is exploited to suppress background from more intense transitions in the cascade. In the summer of 2017, we successfully commissioned a novel target for the  $^{226}\text{Ra}$  charge radius measurement, which is planned to run in 2018. In addition, 2 weeks of beam time were dedicated to observe the 2s-1s transition for the first time, and quantify the background.

WG2 / 79

## Theory of electron scattering and neutrinos

Corresponding Author(s): rocco@ific.uv.es

WG2 / 69

## Theory of neutrino pion production

Corresponding Author(s): raugonjim@gmail.com

WG3 and WG4 - Muon Beam Facilities / 125

## Towards a new High Intensity Muon Beam at PSI: Status and Prospects

Corresponding Author(s): angela.papa@psi.ch

WG3 Parallel Session 3 / 150

## Upgrade Possibility of the ESS Linac for the ESSnuSB Project

Corresponding Author(s): bjorn.galnander@esss.se

WG2 / 132

## **WAGASCI**

**Corresponding Author(s):** p.hallsjo.1@research.gla.ac.uk

**Plenary II / 5**

## **WG1 Goals for the Meeting**

**Corresponding Author(s):** n.mccauley@liv.ac.uk

**Plenary Summary I / 40**

## **WG1 Summary**

**Corresponding Author(s):** bianjm@physics.umn.edu

**Plenary II / 6**

## **WG2 Goals for the Meeting**

**Corresponding Author(s):** kajetan.niewczas@ugent.be

**Plenary Summary I / 41**

## **WG2 Summary**

**Corresponding Author(s):** vishvas@vt.edu

**Plenary II / 7**

## **WG3 Goals for the Meeting**

**Corresponding Author(s):** tetsuro.sekiguchi@kek.jp

**Plenary Summary I / 42**

## **WG3 Summary**

**Corresponding Author(s):** bfreemire@niu.edu

**WG4 -CLFV I / 102**

## **WG4 Charge**

**Plenary II / 8**

## **WG4 Goals for the Meeting**

**Corresponding Author(s):** group@virginia.edu

**Plenary Summary II and Closeout / 37**

## **WG4 Summary**

**Corresponding Author(s):** group@virginia.edu

**WG3 and WG4 - Muon Beam Facilities / 129**

## **WG4 and WG5 - Working Time**

**Plenary II / 9**

## **WG5 Goals for the Meeting**

**Corresponding Author(s):** walter.bonivento@ca.infn.it

**Plenary Summary II and Closeout / 38**

## **WG5 Summary**

**Corresponding Author(s):** walter.bonivento@ca.infn.it

**4**

## **Welcome**

**1**

## **Welcome**

Session 1 / 195

### **Welcome and introduction**

Corresponding Author(s): pahuner@vt.edu

WG4 - CLFV at Colliders / 117

### **Working Group Discussion Time**

WG4 - g-2, precision physics with muons / 112

### **Working Group Discussion Time**

Plenary VI / 30

### **nuSTORM Facility and Report from Cross Section Workshop**

Corresponding Author(s): k.long@imperial.ac.uk

Session 2 / 200

### **nuSTORM accelerator concept to serve cross-section programme**

Corresponding Author(s): sam.tygier@manchester.ac.uk, robert.appleby@manchester.ac.uk, j.pasternak@imperial.ac.uk