




Inelastic neutrino-nucleus scattering in the superscaling model

Jesús González Rosa (jgrosa@us.es)


**Coauthors: Juan Manuel Franco, Juan Antonio Caballero,
Guillermo Daniel Megías, Maria B. Barbaro**

Introduction




- Quasielastic region.
- 2p-2h excitations.
- Δ resonance, other resonances and DIS.

T2K CC ν_μ , $\langle E_{\nu_\mu} \rangle \sim 0.8 \text{ GeV}$, inclusive data
 $0.90 < \cos\theta_\mu < 0.94$




Total Inclusive Cross Section



Superscaling model with QE + 2p2h + 1π [M.V Ivanov et al., J. Phys. G 43, 045101 (2016)].

Introduction, Model: 1π -SuSAv2

Comparison between SuSAv2 and RFG scaling function.



$$f(\psi) = k_F \frac{\left(\frac{d^2\sigma}{d\Omega dw} \right)}{\left(\frac{d^2\sigma}{d\Omega dw} \right)_{s.n}}$$

The scaling function does not depend explicitly on the transferred momentum or the nuclear species [J. E. Amaro et al., J. Phys. G 47, 124001 (2020), G. D. Megias, PhD Thesis (2017)].

SuSAv2 model comes from RMF.

SuSAv2-QE scaling function is going to be implemented in the inelastic regime.

In order to describe the Δ resonance region, a Δ pion production model [PRC 71, 015501 (2005)] is used with the SuSAv2 scaling function.

$$[W^{\mu\nu}]^\Delta = \frac{1}{2} \Lambda_0 f^{model} U^{\mu\nu}$$

Model: SuSAv2-inelastic

SuSAv2-inelastic model describes the full inelastic spectrum (Δ , other res. And DIS) [G. D. Megias, PhD Thesis (2017), M. B. Barbaro et al. Phys. C 69, 035502 (2004)]. Good agreement with (e,e') data.

$$R_{inel}^K(\kappa, \tau) = \frac{N}{\eta_F^2 \kappa} \xi_F \int_{\mu_X^{min}}^{\mu_X^{max}} d\mu_X f^{model}(\psi'_X) U^k$$


The hadronic response is given by an integration of the single-nucleon tensor over the invariant mass.

The limits of this integral are


$$\mu_X^{min} = 1 + \frac{m_\pi}{M_N}, \mu_X^{max} = 1 + 2\lambda - \frac{E_S}{M_N}$$

This limits can be changed to work alongside a resonance model.


BR and BC parametrizations (specially BC) work well. PDF gets closer at high ω , but it is not suited to describe Δ region.



Inelastic Feynmann Diagram



Model: SuSAv2-inelastic



$$F_2^{vN}(x) \approx \frac{18}{5} F_2^{eN}(x)$$

Bodek-Ritchie parametrization
[PRD 23, 1070 (1981)].

Parton Distribution Function.



$$F_3(x) = F_2(x) - 2\bar{Q}(x)$$

Antiquarks
distribution.

$$xF_3^{vp(n)} = 2x(d + s - \bar{u}(\bar{d}) - \bar{c})$$


**Work in
progress for
neutrinos.**

Results



T2K CC ν_{μ} , $\langle E_{\nu_{\mu}} \rangle \sim 0.8 \text{ GeV}$, inclusive data

T2K CC ν_e , $\langle E_{\nu_e} \rangle \sim 1.3 \text{ GeV}$, inclusive data



Total Inclusive Cross Section

Using 1 π – SuSAv2 and the SuSAv2-inelastic model(complete inel). At this kinematics (T2K) DIS does not have a great impact, at higher ω will be more relevant.

Preliminary Result.

Experimental data from [PRL 113(24) (2014)], [PRD 83, 012005 (2011)], [PRD 83, 012005 (2011)] and [PRD 90, 052010 (2014)].

- It is necessary to include an analysis of the inelastic scattering to explain the neutrino cross section at certain kinematics.
- The model works well for electrons and it's expected to work well for neutrinos, specially using BC parametrization in the full inelastic regime.
- In the SuSAv2-inelastic model, the contributions from Δ (π production) and other resonances can be removed , so this model can work alongside other resonance model.
- In the SuSAv2-inelastic model for neutrinos, the analysis of W_3 inelastic structure function is a work in progress.

Thanks for your attention

Limits of the inelastic region



Kinematically
allowed region,
recoiling of the
daughter nucleus

$$\max[\varepsilon(0), 0] \leq \varepsilon \leq \varepsilon(\pi)$$

Considering that the
mass of daughter nuclei
is infinite

$$\varepsilon_{\infty}(\theta) = m_N + \omega - \sqrt{W_X^2 + q^2 + p^2 + 2pq\cos\theta}$$

Considering the
limits

$$m_N + m_{\pi} \leq W_X \leq m_N + \omega - E_S$$

W1, W2 and W3

